

A New IAB Iron Meteorite From China

*K.V. Ponganis, B. Lavielle, B. Spettel, V.F. Buchwald, K.
Nishiizumi, M.W. Caffee, K. Marti*

March 30, 2001

U.S. Department of Energy

Lawrence
Livermore
National
Laboratory

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This report has been reproduced directly from the best available copy.

Available electronically at <http://www.doe.gov/bridge>

Available for a processing fee to U.S. Department of Energy
and its contractors in paper from
U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
Telephone: (865) 576-8401
Facsimile: (865) 576-5728
E-mail: reports@adonis.osti.gov

Available for the sale to the public from
U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
Telephone: (800) 553-6847
Facsimile: (703) 605-6900
E-mail: orders@ntis.fedworld.gov
Online ordering: <http://www.ntis.gov/ordering.htm>

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
<http://www.llnl.gov/tid/Library.html>

A NEW IAB IRON METEORITE FROM CHINA. K.V. Pongani¹, B. Lavielle², B. Spettel³, V.F. Buchwald⁴, K. Nishiizumi⁵, M.W. Caffee⁶, and K. Marti¹, ¹ Dept. of Chemistry & Biochemistry, UCSD, La Jolla, CA 92093-0317, USA (kpongani@ucsd.edu), ²Laboratoire de Chimie Nucléaire Analytique et Bioenvironnementale, U.M.R. 5084 of CNRS, University of Bordeaux I, BP120, Le Haut Vigneau, 33175 Gradignan Cedex, France, ³ Max-Planck-Institute für Chemie, Postfach 3060, D-55020 Mainz, Germany, ⁴ Ingeborgvej 4, DK-2920 Charlottenlund, Denmark, ⁵ Space Sciences Laboratory, UC Berkeley, Berkeley, CA 94720-7450, USA, ⁶ Center for Accelerator Mass Spectrometry, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA.

Introduction: Dr. W. Wei, at UCSD, provided our laboratory with chips from a suspected iron meteorite. The mass (>10 kg) was found approximately 8 km northwest of Zhongshan in Guangxi province by a farmer in his fields. This mass has now been identified as an IAB iron meteorite (provisional name Zhongshan #1) based on its composition, N signature, and petrographic analysis. A second chip from a much larger mass (Zhongshan #2), still in the ground, was recently made available and work is in progress on it, but not reported.

Elemental Composition: A 65 mg sample of Zhongshan #1 was analyzed by INAA at the Max Planck Institute in Mainz. Results are summarized in Table 1. The concentration of diagnostic elements for iron meteorites, such as Ni, Ga, Ge, and platinum group metals, are consistent with the IAB/IIICD group (see Figure 1). [1]




Figure 1. Position of Zhongshan sample #1 in a Ge vs. Ga plot for the IAB/IIICD iron meteorites. Group data is from [1].

Isotopic Signatures of N and Xe: Both N and Xe isotopic signatures were done at UCSD on Zhongshan sample #1. The step-release profile of N is shown in Figure 2. In iron meteorites the signature of indigenous N is highly variable and can be used for classification. [2,3] Nitrogen isotopic analysis indicated that the N step-release profile in this sample is consistent with the IAB/IIICD group, [4] although the total N amounts are extremely small (0.57 ppm). The average $\delta^{15}\text{N}$ for steps 1400 °C to 1600 °C is $-(78 \pm 25) \text{‰}$. There is a different signature for the 900 °C to 1000 °C temperature steps (avg. $\delta^{15}\text{N} = -(40 \pm 7) \text{‰}$), suggesting the presence of at

least two N sources in the sample. The Xe data can be used to trace the presence of inclusions like silicates. [5] The $^{129}\text{Xe}/^{132}\text{Xe}$ ratio (1.38 ± 0.04) shows a ^{129}Xe excess for $1200^\circ\text{C} \leq T \leq 1600^\circ\text{C}$, which is consistent with the presence of minor inclusions.




Figure 2. The stepwise pyrolysis of the Zhongshan iron sample #1. Temperature is in hundreds of °C. The large uncertainties for the steps $\geq 1200^\circ\text{C}$ reflect the low concentrations of N and the variable and large blank corrections.

Cosmogenic Components: An analysis of He, Ne, and Ar on Zhongshan #1 was done at Laboratoire de Chimie Nucléaire Analytique et Bioenvironnementale (see Table 2). The observed ratio $^{36}\text{Ar}/^{38}\text{Ar}$ of 0.6194 represents a spallation component from a shielded location. The $^4\text{He}/^{21}\text{Ne}$ ratio (374 ± 17), the shielding parameter, supports this conclusion.

We can only estimate the cosmic-ray exposure (CRE) age for this meteorite since work on the ^{36}Cl concentration is still in progress. However, assuming a simple exposure history and based on the Ar concentration and on the $^4\text{He}/^{21}\text{Ne}$ ratio, the ^{36}Cl value should be $\leq 11 \text{ dpm/kg}$. (see Figure 3). Using a ^{36}Ar value of $11.38 \times 10^{-8} \text{ cm}^3 \text{ STP/g}$, the CRE age is estimated to be $\geq 440 \text{ Ma}$.

Petrographic Analysis: A sample of Zhongshan #1 was studied at the Geologisk Museum, Copenhagen. A chip was polished, and then, etched lightly with 1% Nitral before analysis.

The bulk of the specimen is kamacite, with a band-width estimated at $(11.9 \pm 0.3) \text{ mm}$. The kamacite, which is rich in Neumann bands, is penetrated by a multitude of shear zones that bend the Neumann bands and shear




Figure 3. Meteorite Shielding: Size vs. Depth. Using the shielding parameter $S = 374$ and the ^{36}Cl profiles from large irons, the expected ^{36}Cl concentration is ≤ 11 dpm/kg. The CRE age is calculated using the following equation: T_c (in Ma) = $427 \times [^{36}\text{Ar}]/[^{36}\text{Cl}]$, where $[^{36}\text{Ar}]$ is in $10^{-8} \text{ cm}^3 \text{ STP/g}$ and $[^{36}\text{Cl}]$ is in dpm/kg. [6] The estimated CRE age is ≥ 440 Ma, based on Old Woman (recovered mass 2753 kg) and Yanhuilitan (recovered mass 421 kg) systematics.

the minerals. The Vickers hardness (200 g load) of the kamacite is: 198 - 199 - 228 - 232 - 233 - 245. These values are high for kamacite, but are due to the significant cold deformations observed in the microstructure.

Taenite is present as up to 0.06 mm wide lamellae, alternating in bluish and brownish colors, suggesting a submicroscopic two-phased structure (probably tetrataenite and associated structures). Three or four plesite fields of the comb plesite variety are present, the largest field being 3×1.5 mm. The bulk Ni (6.6%) seems a little low for the average structure observed.

Schreibersite is abundant as grain boundary crystals (up to 0.08 mm wide). Rhabdites are very common, but most are < 0.02 mm across. Sub-boundaries of the kamacite are decorated with 0.002 mm crystals, probably of phosphides, but possibly supplemented by some carbides. Both schreibersite and rhabdites are severely sheared and displaced. Terrestrial corrosion has had easy access along the grain boundaries with

sheared schreibersite, both because of the capillary action and because of the Ni gradient in kamacite adjacent to schreibersite.

The fusion crust and the heat-affected α_2 zone have been lost by weathering. However, heavy deformation and shearing has generated locally sufficient heat ($> 800^\circ\text{C}$ for a short period of time) to transform kamacite to serrated α_2 grains and remove the Neumann bands. This friction re-heating may have occurred when the meteorite suffered a violent split-up upon entering the Earth's atmosphere. The deformation of the present specimen indicates that it is only a fragment of a bigger shower, consistent with the find nearby of a much larger mass of iron (> 1 ton). The absence of cohenite, graphite, and troilite may be due to the fact that the analysis was performed on a very small, unrepresentative fragment.

Conclusion: Preliminary N isotopic analysis on Zhongshan sample #1 suggested that the iron mass might be a IAB/III CD type. Petrographic inspection and INAA analysis has verified that this sample is a typical member of the coarse octahedrites, of the resolved chemical group IAB. Zhongshan #1 was well-shielded (see Figure 3). Its estimated CRE age is ≥ 440 Ma. Determination of the radionuclides and noble gases on Zhongshan sample #2 (in progress) will give useful information on the exposure and shielding history of the meteoroid and its possible atmospheric break-up.

Acknowledgements: We are grateful to Dr. W. Wei at UCSD for bringing the meteorite to our attention and to Dr. J. Zipfel at Max-Planck-Institut für Chemie for her help in this work.

References: [1] Choi, B.G., *et al.*, (1995) *Geochimica Et Cosmochimica Acta* 59: 593-612. [2] Prombo, C.A. and Clayton, R.N., (1993) *Geochimica et Cosmochimica Acta* 57: 3749-3761. [3] Franchi, I.A., *et al.*, (1993) *Geochimica et Cosmochimica Acta* 57: 3105-3121. [4] Ponganis, K.V., *et al.*, (2000) *Meteoritics and Planetary Science* 35 (Supplement): A128-129. [5] Ponganis, K.V. and Marti, K., (2000) *LPSC XXXI*: [6] Lavielle, B., *et al.*, (1999) *Earth and Planetary Science Letters* 170: 93-104.

Table 1. Trace elements of Zhongshan #1 sample. Fe is 91.94% (3% error) and Ni is 6.60% (4% error), by mass.

| Element | ppm | % error | Element | ppm | % error | Element | ppm | % error |
|---------|-------|---------|---------|-------|---------|---------|-------|---------|
| Co | 4330 | 3 | Mo | 8.20 | 7 | Re | 0.200 | 5 |
| Cu | 136 | 4 | Ru | 5.70 | 15 | Os | 2.90 | 7 |
| Ga | 88.6 | 3 | Pd | 3.60 | 15 | Ir | 2.34 | 3 |
| Ge | 301 | 7 | Sb | 0.390 | 10 | Pt | 7.20 | 7 |
| As | 11.9 | 4 | W | 1.01 | 5 | Au | 2.22 | 3 |
| Se | < 4 | - | | | | | | |

Table 2. Cosmogenic noble gas concentrations (in $10^{-8} \text{ cm}^3 \text{ STP/g}$) and ratios for the Zhongshan #1 sample.

| | ^4He | ^{21}Ne | ^{36}Ar | ^{38}Ar | $^3\text{He}/^4\text{He}$ | $^{36}\text{Ar}/^{38}\text{Ar}$ | $^4\text{He}/^{21}\text{Ne}$ | $^4\text{He}/^{38}\text{Ar}$ | $^{21}\text{Ne}/^{38}\text{Ar}$ |
|---------------------|---------------|------------------|------------------|------------------|---------------------------|---------------------------------|------------------------------|------------------------------|---------------------------------|
| value | 1120 | 2.995 | 11.38 | 18.37 | 0.2435 | 0.6194 | 374 | 61.0 | 0.1631 |
| error (2 σ) | 37 | 0.093 | 0.19 | 0.29 | 0.0040 | 0.0037 | 17 | 2.3 | 0.0057 |